

Melville (pers. comm.) also agreed, stating, however, that *goisagi* has blue facial skin when breeding (shown by one individual in Hong Kong), and greenish-yellow the rest of the year. He suggested that the pale spotting on the neck indicated that it was an immature *goisagi*.

Wells (pers. comm. to C. F. Mann and the author) thought that the pattern of the underparts could be matched by immature *melanolophus*, but the pale, indistinctly vermiculated upperparts were wrong for that species. He felt that the bare parts coloration was inconclusive but agreed that bird must have been *goisagi*.

The species breeds in southern Japan, migrating to eastern China, Taiwan, the Philippines and Sulawesi; vagrants have reached Sakhalin, Hokkaido and Korea.

I would like to thank C. F. Mann for help in preparing this note.

REFERENCE

Mann, C. F. (1988) Bird report for Brunei Darussalam 1986-88. *Brunei Mus. J.* 6(4): 88-111.

Black-collared Starling *Sturnus nigricollis*, a species new to Borneo and the Sundas

JEN A. ELKIN, CLIVE F. MANN and A. CONRAD OZOG

At about 08h00 on 4 October 1991 a large starling was seen flying about 10 m above a fallow paddyfield at Wasan Rice Scheme, Brunei. Its large size (almost that of a Hill Myna *Gracula religiosa*) and pied appearance indicated that it was not one of the starling species regularly occurring in Borneo. It dropped on a path about 300 m away and was eventually relocated with flocks of egrets *Egretta* and *Bubulcus* and terns *Chlidonias* following a plough. It was observed at close quarters with both binoculars and a telescope and the following points were noted:

Size of Common Myna *Acridotheres tristis*, or slightly larger. Bill and legs pale. Back, wings, mantle and tail blackish-brown or black; a white bar on primaries, particularly noticeable in flight; rump and tips of tail feathers white. Head, neck and chin to breast greyish-brown to buffish-brown, streaked whitish and palest on the throat.

It was identified as an immature Black-collared Starling *Sturnus nigricollis* from King *et al.* (1975) and Viney and Phillipps (1988). The species occurs from southern China south to Myanmar and Thailand but has not previously been recorded in the Sundas.

REFERENCES

- King, B. F., Dickinson, E. C. and Woodcock, M. W. (1975) *A field guide to the birds of South-East Asia*. London: Collins.
 Viney, C. and Phillipps, K. (1988) *New colour guide to Hong Kong birds*. Fourth edition. Hong Kong: Government Printer.

Jen Elkin, PTR/3, Brunei Shell Pteroleum, Seria, Brunei

Clive F. Mann, 123 Hartwood Road, London W12 9NG, U.K.

A. Conrad Ozog, c/o UBD, PO Gadong, Bandar Seri Begawan 3186, Brunei.

Long-billed Partridge *Rhizothera longirostris*: a new species for Sabah

D. A. SHOWLER

The Long-billed Partridge *Rhizothera longirostris* is represented in Borneo by two races, neither of which has been recorded previously in Sabah (Smythies 1981, Francis 1986). *R. l. dulitensis* apparently occurs only on Mt. Dulit and Mt. Batu Song in Sarawak, above 1,000 m. *R. l. longirostris* is the lowland race, known previously only from south-west Sarawak and the Barito drainage in south Kalimantan (Smythies 1981). It is presumably this latter race that has been seen and heard by several observers in primary lowland dipterocarp rainforest at Danum valley, east Sabah. It was apparently first recorded in the area by R. Orenstein (Toronto, Canada) and two Australian birders, during the third week of March 1992. I heard a pair duetting at dusk on 27 March 1992, 500 m along the main trail west (W5), and about 1 km from Danum Valley Field Centre. I subsequently heard a pair (probably the same birds) duetting, and saw a male walking back into cover, at 17h15 on 4 April 1992, 300-400 m along the main trail west (W3-W4).

REFERENCES

- Francis, C. M. (1986) *Check-list of the birds of Sabah*.
 Smythies, B. E. (1981) *The birds of Borneo*. Third edition. Kota Kinabalu and Kuala Lumpur: The Sabah Society and the Malayan Nature Society.

D. A. Showler, 30 Essex Street, Norwich, NR2 2BL, U.K.